## Uncertainties in Aspheric Profile Measurements with the Geometry Measuring Machine at NIST

B. Chang Kim<sup>1</sup>, Nadia Machkour-Deshayes<sup>1</sup>, Quandou Wang<sup>1</sup>, Johannes Soons<sup>1</sup>,Ulf Griesmann<sup>1</sup>, Lahsen Assoufid<sup>2</sup>

<sup>1</sup>National Institute of Standards & Technology, Dimensional Metrology Program, Gaithersburg, Maryland 20899 <sup>2</sup> XFD, Argonne National Laboratory, 9700 South Cass Avenue, Argonne, IL 60439

The Geometry Measuring Machine (GEMM) at the National Institute of Standards and Technology (NIST) is a profilometer for free-form surfaces. It measures the local curvature of a test part surface at several locations along a line. The profile is then reconstructed from the curvature data. For form measurements of free-form surfaces, methods based on local part curvature sensing have strong appeal. Unlike full-aperture interferometry they do not require customized null optics. The uncertainty of the reconstructed profile is critically dependent on the curvature measurement uncertainty. We evaluate the measurement uncertainties for a curvature sensor based on a small aperture interferometer and the resulting uncertainty for the reconstructed profile. Profile measurements of a free-form mirror made with GEMM are compared with measurements using a long-trace profiler and subaperture-stitching interferometry.